

## NEAREST POINTS AND DELTA CONVEX FUNCTIONS IN BANACH SPACES

JONATHAN M. BORWEIN and OHAD GILADI

### Abstract

Given a closed set  $C$  in a Banach space  $(X, \|\cdot\|)$ , a point  $x \in X$  is said to have a nearest point in  $C$  if there exists  $z \in C$  such that  $d_C(x) = \|x - z\|$ , where  $d_C$  is the distance of  $x$  from  $C$ . We shortly survey the problem of studying the size of the set of points in  $X$  which have nearest points in  $C$ . We then turn to the topic of delta-convex functions and indicate how it is related to finding nearest points.

2010 *Mathematics subject classification*: primary 46B10; secondary 41A29.

*Keywords and phrases*: Nearest points, Fréchet sub-differentials, delta convex functions.

### 1. Nearest points in Banach spaces

**1.1. Background** Let  $(X, \|\cdot\|)$  be a real Banach space, and let  $C \subseteq X$  be a non-empty closed set. Given  $x \in X$ , its distance from  $C$  is given by

$$d_C(x) = \inf_{y \in C} \|x - y\|.$$

If there exists  $z \in C$  with  $d_C(x) = \|x - z\|$ , we say that  $x$  has a *nearest point* in  $C$ . Let also

$$N(C) = \{x \in X : x \text{ has a nearest point in } C\}.$$

One can then ask questions about the structure of the set  $N(C)$ . This question has been studied in [4, 11, 14, 20, 22, 27, 28, 30, 35] to name just a few articles. More specifically, the following questions are at the heart of this note:

*Given a nonempty closed set  $C \subseteq X$ , how large is the set  $N(C)$ ? When is  $N(C)$  non-empty?*

One way to answer this question is to consider sets which are large in the set-theoretic topological sense, such as dense  $G_\delta$  sets. We begin with a few definitions.

**DEFINITION 1.1.** If  $N(C) = X$ , i.e., if every point in  $X$  has a nearest point in  $C$ , then  $C$  is said to be proximal. If  $N(C)$  contains a dense  $G_\delta$  set, then  $C$  is said to be almost proximal.

In passing, we recall that if every point in  $X$  is uniquely proximal then  $C$  is said to be a *Chebyshev set*. It has been conjectured for over half a century, that in Hilbert space Chebyshev sets are necessarily convex, but this is only proven for weakly closed sets [7]. See also [17] for a recent survey of this topic that, in particular, give a clear construction of a non-convex Chebyshev set in an incomplete inner product space.

For example, closed convex sets in reflexive spaces are proximal, as are all closed sets in finite dimensional spaces. See [4]. One can also consider stronger notions of "large" sets. See Section 1.4. First, we also need the following concept.

**DEFINITION 1.2.** A Banach space is said to be a (sequentially) Kadec space if for each sequence  $\{x_n\}$  that converges weakly to  $x$  with  $\lim \|x_n\| = \|x\|$ ,  $\{x_n\}$  converges to  $x$  in norm, i.e.,

$$\lim_{n \rightarrow \infty} \|x - x_n\| = 0.$$

All locally uniformly convex Banach spaces are Kadec spaces as are all finite dimensional spaces. With the above definitions in hand, the following lovely result holds.

**THEOREM 1.3** (Lau [22], Borwein-Fitzpatrick [4]). *If  $X$  is a reflexive Kadec space and  $C \subseteq X$  is closed, then  $C$  is almost proximal.*

The assumptions on  $X$  are in fact necessary.

**THEOREM 1.4** (Konjagin [20]). *If  $X$  is not both Kadec and reflexive, then there exist  $C \subseteq X$  closed and  $U \subseteq X \setminus C$  open such that no  $x \in U$  has a nearest point in  $C$ .*

It is known that under stronger assumption on  $X$  one can obtain stronger results on the set  $N(C)$ . See Section 1.4.

## 1.2. Fréchet sub-differentiability and nearest points

We begin with a definition.

**DEFINITION 1.5.** Assume that  $f : X \rightarrow \mathbb{R}$  is a real valued function with  $f(x)$  finite. Then  $f$  is said to be Fréchet sub-differentiable at  $x \in X$  if there exists  $x^* \in X^*$  such that

$$\liminf_{y \rightarrow 0} \frac{f(x+y) - f(x) - x^*(y)}{\|y\|} \geq 0. \quad (1.1)$$

The set of points in  $X^*$  that satisfy (1.1) is denoted by  $\partial f(x)$ .

Sub-derivatives have been found to have many applications in approximation theory. See for example [4, 5, 7, 8, 25].

One of the connections between sub-differentiability and the nearest point problem was studied in [4]. Given  $C \subseteq X$  closed, the following modification of a construction of [22] was introduced. Consider

$$L_n(C) = \left\{ x \in X \setminus C : \exists x^* \in \mathbb{S}_{X^*} \text{ s.t. } \sup_{\delta > 0} \inf_{z \in C \cap B(x, d_C(x) + \delta)} x^*(x - z) > (1 - 2^{-n})d_C(x) \right\},$$

where  $\mathbb{S}_{X^*}$  denotes the unit sphere of  $X^*$ . Also, let

$$L(C) = \bigcap_{n=1}^{\infty} L_n(C).$$

The following is known.

**PROPOSITION 1.6 (Borwein-Fitzpatrick [4]).** *For every  $n \in \mathbb{N}$ ,  $L_n(C)$  is open. In particular,  $L(C)$  is  $G_\delta$ .*

Finally, let

$$\Omega(C) = \left\{ x \in X \setminus C : \exists x^* \in \mathbb{S}_{X^*}, \text{ s.t. } \forall \epsilon > 0, \exists \delta > 0, \right. \\ \left. \inf_{z \in C \cap B(x, d_C(x) + \delta)} x^*(x - z) > (1 - \epsilon)d_C(x) \right\}.$$

While  $L(C)$  is  $G_\delta$  by Proposition 1.6, under the assumption that  $X$  is reflexive, the following is known.

**PROPOSITION 1.7 (Borwein-Fitzpatrick [4]).** *If  $X$  is reflexive then  $\Omega(C) = L(C)$ . In particular,  $\Omega(C)$  is  $G_\delta$ .*

The connection to sub-differentiability is given in the following proposition.

**PROPOSITION 1.8 (Borwein-Fitzpatrick [4]).** *If  $x \in X \setminus C$  and  $\partial d_C(x) \neq \emptyset$ , then  $x \in \Omega(C)$ .*

The following fundamental result is available.

**THEOREM 1.9 (Borwein-Preiss [6]).** *If  $f$  is lower semicontinuous on a reflexive Banach space, then  $f$  is Fréchet sub-differentiable on a dense set.*

In fact, Theorem 1.9 holds under a weaker assumption. See [4, 6]. Since the distance function is lower semicontinuous, it follows that it is sub-differentiable on a dense subset, and therefore, by the above propositions,  $\Omega(C)$  is a dense  $G_\delta$  set. Thus, in order to prove Theorem 1.3, it is only left to show that every  $x \in \Omega(C)$  has a nearest point in  $C$ . Indeed, if  $\{z_n\} \subseteq C$  is a minimizing sequence, then by extracting a subsequence, assume that  $\{z_n\}$  has a weak limit  $z \in C$ . By the definition of  $\Omega(C)$ , there exists  $x^* \in \mathbb{S}_{X^*}$  such that

$$\|x - z\| \geq x^*(x - z) = \lim_{n \rightarrow \infty} x^*(x - z_n) \geq d_C(x) = \lim_{n \rightarrow \infty} \|x - z_n\|.$$

On the other hand, by weak lower semicontinuity of the norm,

$$\lim_{n \rightarrow \infty} \|x - z_n\| \geq \|x - z\|,$$

and so  $\|x - z\| = \lim \|x - z_n\|$ . Since it is known that  $\{z_n\}$  converges weakly to  $z$ , the Kadec property implies that in fact  $\{z_n\}$  converges in norm to  $z$ . Thus  $z$  is a nearest point. This completes the proof of Theorem 1.3.

This scheme of proof, taken from [4], shows that differentiation arguments can be fruitfully used to prove that  $N(C)$  is large.

**1.3. Nearest points in non-Kadec spaces** It was previously mentioned that closed convex sets in reflexive spaces are proximal. It is also known that non-empty “Swiss cheese” sets (sets whose complement is a mutually disjoint union of open convex sets) in reflexive spaces are almost proximal [4]. These two examples show that for some classes of closed sets, the Kadec property can be removed. Moreover, one can consider another, weaker, way to “measure” whether a set  $C \subseteq X$  has “many” nearest points: ask whether the set of nearest points in  $C$  to points in  $X \setminus C$  is dense in the boundary of  $C$ . Note that if  $C$  is almost proximal, then nearest points are dense in the boundary. The converse, however, is not true. In [4] an example of a non-Kadec reflexive space was constructed where for every closed set, the set of nearest points is dense in its boundary. The following general question is still open, even in renormings of Hilbert space.

**QUESTION 1.10.** *Let  $(X, \|\cdot\|)$  be a reflexive Banach space and suppose  $C \subseteq X$  closed. Is the set of nearest points in  $C$  to points in  $X \setminus C$  dense in its boundary?*

Relatedly, if the set  $C$  is norm closed and bounded in a space with the *Radon-Nikodym property* as is the case of reflexive space, then  $N(C)$  is nonempty and is large enough so that  $\overline{\text{conv}}C = \overline{\text{conv}}N(C)$  [4].

**1.4. Porosity and nearest points** As was mentioned in subsection 1.2, one can consider stronger notions of “large” sets. One is the following notion.

**DEFINITION 1.11.** A set  $S \subseteq X$  is said to be porous if there exists  $c \in (0, 1)$  such that for every  $x \in X$  and every  $\epsilon > 0$ , there is a  $y \in B(0, \epsilon) \setminus \{0\}$  such that

$$B(x + y, c\|y\|) \cap S = \emptyset.$$

A set is said to be  $\sigma$ -porous if it is a countable union of porous sets. Here and in what follows,  $B(x, r)$  denotes the closed ball around  $x$  with radius  $r$ .

It is known that every  $\sigma$ -porous set is of the first category, i.e., union of nowhere dense sets. Moreover, it is also known that the class of  $\sigma$ -porous sets is a proper sub-class of the class of first category sets. When  $X = \mathbb{R}^n$ , one can show that every  $\sigma$ -porous set has Lebesgue measure zero. This is not the case for every first category set:  $\mathbb{R}$  can be written as a disjoint union of a set of the first category and a set of Lebesgue measure zero. Hence, the notion of porosity automatically gives a stronger notion of large sets: every set whose complement is  $\sigma$ -porous is also a dense  $G_\delta$  set. We recommend [23, 36] for a more detailed discussion on porous sets.

We recall that a Banach space  $(X, \|\cdot\|)$  is said to be *uniformly convex* if the function

$$\delta(\epsilon) = \inf \left\{ 1 - \left\| \frac{x+y}{2} \right\| : x, y \in \mathbb{S}_X, \|x-y\| \geq \epsilon \right\}, \quad (1.2)$$

is strictly positive whenever  $\epsilon > 0$ . Here  $\mathbb{S}_X$  denotes the unit sphere of  $X$ . In [11] the following was shown.

**THEOREM 1.12** (De Blasi-Myjak-Papini [11]). *If  $X$  is uniformly convex, then  $N(C)$  has a  $\sigma$ -porous complement.*

In fact, [11] proved a stronger result, namely that for every  $x$  outside a  $\sigma$ -porous set, the minimization problem is *well posed*, i.e., there is unique minimizer to which every minimizing sequence converges. See also [16, 27, 28] for closely related results in this direction.

The proof of Theorem 1.12 builds on ideas developed in [30]. It would, however, be interesting to know whether one may use differentiation arguments as in Section 1.2. This raises the following question:

**QUESTION 1.13.** *Can differentiation arguments be used to give an alternative proof of Theorem 1.12?*

More specifically, if one can show that  $\partial d_C \neq \emptyset$  outside a  $\sigma$ -porous set, then by the arguments presented in Section 1.2, it would follow that  $N(C)$  has a  $\sigma$ -porous complement. Next, we mention two important results regarding differentiation in Banach spaces. See also [23, Sec. 3.3].

**THEOREM 1.14** (Preiss-Zajíček [26]). *If  $X$  has a separable dual and  $f : X \rightarrow \mathbb{R}$  is continuous and convex, then  $X$  is Fréchet differentiable outside a  $\sigma$ -porous set.*

Theorem 1.14 implies that if, for example,  $d_C$  is a linear combination of convex functions (see more on this in Section 2), then  $N(C)$  has a  $\sigma$ -porous complement. Also, we have the following.

**THEOREM 1.15** (Cúth-Rmoutil [10]). *If  $X$  has a separable dual and  $f : X \rightarrow \mathbb{R}$  is Lipschitz, then the set of points where  $f$  is Fréchet sub-differentiable but not differentiable is  $\sigma$ -porous.*

Since  $d_C$  is 1-Lipschitz (non-expansive), the issues from a porosity perspective of seeking points of sub-differentiability or points of differentiability are similar. We also observe that Theorem 1.14 and Theorem 1.15 remain true if we consider  $f : A \rightarrow \mathbb{R}$  where  $A \subseteq X$  is open and convex.

We now turn from results depending on the geometry of the space to those exploiting the finer structure of  $D_C$  or  $C$ .

## 2. DC functions and DC sets

### 2.1. Background

**DEFINITION 2.1.** A function  $f : X \rightarrow \mathbb{R}$  is said to be delta-convex, or DC, if it can be written as a difference of two convex functions on  $X$ .

This notion was introduced in [18] and was later studied by many authors. See, for example, [2, 8, 9, 12, 13, 21, 24, 34]. In particular, [2] gives a concise introduction

to this topic. We will discuss here only the parts that are closely related to the nearest point problem.

The following is an important and attractive proposition. See, for example, [19, 33] for a proof.

**PROPOSITION 2.2.** *If  $f_1, \dots, f_k$  are DC functions and  $f : X \rightarrow \mathbb{R}$  is continuous and  $f(x) \in \{f_1(x), \dots, f_n(x)\}$ . Then  $f$  is also DC.*

The result remains true if we replace the domain  $X$  by any convex subset.

**2.2. DC functions and nearest points** Showing that a given function is in fact DC is a powerful tool, as it allows us to use many known results about convex and DC functions. For example, if a function is DC on a Banach space with a separable dual, then by Theorem 1.14, it is differentiable outside a  $\sigma$ -porous set. In the context of the nearest point problem, if we know that the distance function is DC, then using the scheme presented in Section 1.2, it follows that  $N(C)$  has a  $\sigma$ -porous complement. The same holds if we have a difference of a convex function and, say, a smooth function.

The simplest and best known example, originally due to Asplund, is that when  $(X, \|\cdot\|)$  is a Hilbert space, we have the following:

$$\begin{aligned} d_C^2(x) &= \inf_{y \in C} \|x - y\|^2 \\ &= \inf_{y \in C} [\|x\|^2 - 2\langle x, y \rangle + \|y\|^2] \\ &= \|x\|^2 - 2 \sup_{y \in C} [\langle x, y \rangle - \|y\|^2/2], \end{aligned}$$

and the function  $x \mapsto \sup_{y \in C} [\langle x, y \rangle - \|y\|^2/2]$  is convex as a supremum of affine functions. Hence  $d_C^2$  is DC on  $X$ . Moreover, in a Hilbert space we have the following result (see [8, Sec. 5.3]).

**THEOREM 2.3.** *If  $(X, \|\cdot\|)$  is a Hilbert space,  $d_C$  is locally DC on  $X \setminus C$ .*

**PROOF.** Fix  $y \in C$  and  $x_0 \in X \setminus C$ . It can be shown that if we let  $f_y(x) = \|x - y\|$ , then  $f_y$  satisfies

$$\|f'_y(x_1) - f'_y(x_2)\|_{X^*} \leq L_{x_0} \|x_1 - x_2\|, \quad x_1, x_2 \in B_{x_0},$$

where  $L_{x_0} = \frac{4}{d_S(x_0)}$  and  $B_{x_0} = B(x_0, \frac{1}{2}d_C(x_0))$ . In particular,

$$(f'_y(x + t_1 v) - f'_y(x + t_2 v))(v) \leq L_{x_0}(t_2 - t_1), \quad v \in \mathbb{S}_X, t_2 > t_1 \geq 0, \quad (2.1)$$

whenever  $x + t_1 v, x + t_2 v \in B_{x_0}$ . Next, the convex function  $F(x) = \frac{L_{x_0}}{2} \|x\|^2$  satisfies

$$(F'(x_1) - F'(x_2))(x_1 - x_2) \geq L_{x_0} \|x_1 - x_2\|^2, \quad \forall x_1, x_2 \in X. \quad (2.2)$$

In particular

$$(F'(x + t_2v) - F'(x + t_1v))(v) \geq L_{x_0}(t_2 - t_1), \quad v \in \mathbb{S}_X, \quad t_2 > t_1 \geq 0. \quad (2.3)$$

Altogether, if  $g_y(x) = F(x) - f_y(x)$ , then

$$(g'_y(x + t_2v) - g'_y(x + t_1v))(v) \stackrel{(2.1) \wedge (2.3)}{\geq} 0, \quad v \in \mathbb{S}_X, \quad t_2 > t_1 \geq 0,$$

whenever  $x + t_1v, x + t_2v \in B_{x_0}$ . This monotonicity implies that  $g_y$  is convex on  $B_{x_0}$  [7]. It then follows that

$$d_C(x) = \frac{L_{x_0}}{2} \|x\|^2 - \sup_{y \in C} \left[ \frac{L_{x_0}}{2} \|x\|^2 - \|x - y\| \right] = h(x) - \sup_{y \in C} g_y(x)$$

is DC on  $B_{x_0}$ . □

**REMARK 2.4.** *Even in  $\mathbb{R}^2$  there are sets for which  $d_C$  is not DC everywhere (not even locally DC), as was shown in [2]. Thus, the most one could hope for in Theorem 2.3 is a locally DC function on  $X \setminus C$ .*

Given  $q \in (0, 1]$ , a norm  $\|\cdot\|$  is said to be *q-Hölder smooth* at a point  $x \in X$  if there exists a constant  $K_x \in (0, \infty)$  such that for every  $y \in \mathbb{S}_X$  and every  $\tau > 0$ ,

$$\frac{\|x + \tau y\|}{2} + \frac{\|x - \tau y\|}{2} \leq 1 + K_x \tau^{1+q}.$$

If  $q = 1$  then  $(X, \|\cdot\|)$  is said to be *Lipschitz smooth* at  $x$ . The spaces  $L_p$ ,  $p \geq 2$  are known to be Lipschitz smooth, and in general  $L_p$ ,  $p > 1$ , is *s-Hölder smooth* with  $s = \min\{1, p - 1\}$ .

A Banach space is said to be *p-uniformly convex* if for every  $x, y \in \mathbb{S}_X$ ,

$$1 - \left\| \frac{x + y}{2} \right\| \geq L \|x - y\|^p.$$

Note that this is similar to assuming that  $\delta(\epsilon) = L\epsilon^p$  in (1.2). The spaces  $L_p$ ,  $p > 1$ , are *r-uniformly convex* with  $r = \max\{2, p\}$ .

One might well ask whether the scheme of proof of Theorem 2.3 can be used in a more general setting. The results which follow indicate that this is not possible.

**PROPOSITION 2.5.** *Let  $(X, \|\cdot\|)$  be a Banach space,  $C \subseteq X$  a closed set, and fix  $x_0 \in X \setminus C$  and  $y \in C$ . Assume that there exists  $r_0$  such that  $f_y(x) = \|x - y\|$  has a Lipschitz derivative on  $B(x_0, r_0)$ :*

$$\|f'_y(x_1) - f'_y(x_2)\| \leq L_{x_0} \|x_1 - x_2\|. \quad (2.4)$$

*Then the norm is Lipschitz smooth on  $-y + B_{x_0} = B(x_0 - y, r_0)$ . If in addition there exists a function  $F : X \rightarrow \mathbb{R}$  satisfying*

$$(F'(x_1) - F'(x_2))(x_1 - x_2) \geq L_{x_0} \|x_1 - x_2\|^2, \quad \forall x_1, x_2 \in B(x_0, r_0), \quad (2.5)$$

*then  $(X, \|\cdot\|)$  admits an equivalent norm which is 2-uniformly convex. In particular, if  $X = L_p$  then  $p = 2$ .*

**PROOF.** To prove the first assertion note that (2.4) is equivalent to

$$\|x - y + h\| + \|x - y - h\| - 2\|x - y\| \leq L_{x_0}\|h\|^2, \quad x \in B_{x_0}.$$

See for example [15, Prop. 2.1].

To prove the second assertion, note that a function that satisfies (2.5) is also known as *strongly convex*: one can show that (2.5) is in fact equivalent to the condition

$$f\left(\frac{x_1 + x_2}{2}\right) \leq \frac{1}{2}f(x_1) + \frac{1}{2}f(x_2) - C\|x_1 - x_2\|^2,$$

for some constant  $C$ . See for example [29, App. A]. This implies that there exists an equivalent norm which is 2-uniformly convex ([7, Thm 5.4.3]).

□

**REMARK 2.6.** From [1] it is known that if  $F : X \rightarrow \mathbb{R}$  satisfies

$$(F'(x_1) - F'(x_2))(v) \geq L\|x_1 - x_2\|^2,$$

for all  $x_1, x_2 \in X$ , and also that  $F$  is twice (Fréchet) differentiable at one point, then  $(X, \|\cdot\|)$  is isomorphic to a Hilbert space.

**REMARK 2.7.** If we replace the Lipschitz condition by a Hölder condition

$$\|f'_y(x_1) - f'_y(x_2)\| \leq \|x_1 - x_2\|^\beta, \quad \beta < 1,$$

then in order to follow the same scheme of proof of Theorem 2.3, instead of (2.2), we would need a function  $F$  satisfying

$$(F'(x_1) - F'(x_2))(x_1 - x_2) \geq \|x_1 - x_2\|^{1+\beta}, \quad x_1, x_2 \in B_{x_0}.$$

which implies

$$\|F'(x_1) - F'(x_2)\| \geq \|x_1 - x_2\|^\beta, \quad x_1, x_2 \in B_{x_0}. \quad (2.6)$$

If  $G = (F')^{-1}$ , then we get

$$\|Gx_1 - Gx_2\| \leq \|x_1 - x_2\|^{1/\beta}, \quad x_1, x_2 \in F'(B_{x_0}),$$

which can occur only if  $G$  is a constant. Hence (2.6) cannot hold and the scheme of proof cannot be generalized if we replace the Lipschitz condition by a Hölder condition.

**2.3. DC sets, DC representable sets** The next definition is quite natural.

**DEFINITION 2.8.** A set  $C$  is said to be a DC set if  $C = A \setminus B$  where  $A, B$  are convex.

We can also consider the following class of sets.

**DEFINITION 2.9.** A set  $C \subseteq X$  is said to be DC representable if there exists a DC function  $f : X \rightarrow \mathbb{R}$  such that  $C = \{x \in X : f(x) \leq 0\}$ .

Note that if  $C = A \setminus B$  is a DC set, then we can write  $C = \{\mathbb{1}_B - \mathbb{1}_A + 1/2 \leq 0\}$ , where  $\mathbb{1}_A, \mathbb{1}_B$  are the convex indicator functions of  $A, B$ , respectively. Therefore,  $C$  is DC representable. Moreover, we have the following.

**THEOREM 2.10 (Thach [31]).** *Assume that  $X$  and  $Y$  are two Banach spaces, and  $T : Y \rightarrow X$  a surjective bounded linear map which is not an isomorphism, i.e.,  $\ker(T) \neq \{0\}$ . Then for any set  $M \subseteq X$  there exists a DC representable set  $D \subseteq Y$ , such that  $M = T(D)$ .*

Also, the following ‘converse’ is known. See [19].

**PROPOSITION 2.11.** *If  $C$  is a DC representable set, then there exist  $A, B \subseteq X \oplus \mathbb{R}$  convex, such that  $x \in C \iff (x, x') \in A \setminus B$ .*

**PROOF.** Define  $g_1(x, x') = f_1(x) - x', g_2(x, x') = f_2(x) - x'$ . Let  $A = \{(x, x') : g_1(x, x') \leq 0\}, B = \{(x, x') : g_2(x, x') \leq 0\}$ . Then  $x \in C \iff (x, x') \in A \setminus B$ .  $\square$

In particular, every DC representable set in  $X$  is a projection of a DC set in  $X \oplus \mathbb{R}$ . The next theorem was proved in [32].

**THEOREM 2.12 (Thach-Konno [32]).** *If  $X$  is a reflexive Banach space and  $C \subseteq X$  is closed, then  $C$  is DC representable.*

This makes relevant the following question.

**QUESTION 2.13.** *Are there any classes of spaces  $X$ , say uniformly convex spaces, such that there exists  $\alpha > 0$  such that  $d_C^\alpha$  is locally DC on  $X \setminus C$  whenever  $C$  is a DC representable set?*

If the answer to Question 2.13 is positive, then by the discussion in subsection 1.2 we can conclude that  $N(C)$  has a  $\sigma$ -porous complement, thus giving an alternative proof of Theorem 1.12. One may also ask Question 2.13 for DC sets instead of DC representable sets.

To end this note, we discuss some simple cases where DC and DC representable sets can be used to study the nearest point problem.

**PROPOSITION 2.14.** *Assume that  $C = X \setminus \bigcup_{a \in \Lambda} U_a$ , where each  $U_a$  is an open convex set. Then  $d_C$  is locally DC (in fact, locally concave) on  $X \setminus C$ .*

**PROOF.** First, it is shown in [4, Sec. 3] that if  $a \in \Lambda$ , then  $d_{X \setminus U_a}$  is concave on  $U_a$ . Next, it also shown in [4] that if  $x \in U_a$  then  $d_{X \setminus U_a}(x) = d_C(x)$ . In particular,  $d_C$  is concave on  $U_a$ .  $\square$

**PROPOSITION 2.15.** *Assume that  $C = A \setminus B$  is a closed DC set, and assume  $A$  is closed and  $B$  is open, then  $d_C$  is convex whenever  $d_C(x) \leq d_{A \cap B}$ .*

**PROOF.** Since  $A = (A \setminus B) \cup B$ , we have

$$d_A(x) = \min \{d_{A \setminus B}(x), d_{A \cap B}(x)\} = \min \{d_C(x), d_{A \cap B}(x)\}.$$

Hence, if  $d_C(x) \leq d_{A \cap B}(x)$  then  $d_C(x) = d_A(x)$  is convex.  $\square$

**PROPOSITION 2.16.** *Assume that  $C$  is a DC representable set, i.e.,  $C = \{x \in X : f_1(x) - f_2(x) \leq 0\}$ , and that  $f_2(x) = \max_{1 \leq i \leq m} \varphi_i(x)$ , where  $\varphi_i$  is affine. Then  $d_C$  is DC on  $X$ .*

**PROOF.** Write

$$\begin{aligned} C &= \{x : f_1(x) - f_2(x) \leq 0\} \\ &= \{x : f_1(x) - \max_{1 \leq i \leq m} \varphi_i(x) \leq 0\} \\ &= \{x : \min_{1 \leq i \leq m} (f_1(x) - \varphi_i(x)) \leq 0\} \\ &= \bigcup_{i=1}^m \{x : f_1(x) - \varphi_i(x) \leq 0\}. \end{aligned}$$

where the sets  $\{x : f_1(x) - \varphi_i(x) \leq 0\}$  are convex sets. Hence, we have that

$$d_C(x) = \min_{1 \leq i \leq m} d_{C_i}(x),$$

is a minimum of convex sets and therefore by Proposition 2.2 is a DC function.  $\square$

In [9] it was shown that if  $X$  is superreflexive, then any Lipschitz map is a uniform limit of DC functions. See also [7, Sec. 5.1]. We have the following simple partner result.

**PROPOSITION 2.17.** *If  $X$  is separable, then  $d_C$  is a limit (not necessarily uniform) of DC functions.*

**PROOF.** If  $X$  is separable, i.e., there exists a countable  $Q = \{q_1, q_2, \dots\} \subseteq X$  with  $\bar{Q} = X$ . We have

$$d_C(x) = \inf_{z \in C} \|x - z\| = \inf_{z \in C \cap Q} \|x - z\| = \lim_{n \rightarrow \infty} \left[ \min_{z \in C \cap Q_n} \|x - z\| \right],$$

where  $Q_n = \{q_1, q_2, \dots, q_n\}$ . Again by Proposition 2.2 we have that  $\min_{z \in C \cap Q_n} \|x - z\|$  is a DC function as a minimum of convex functions.  $\square$

### 3. Conclusion

Despite many decades of study, the core questions addressed in this note are still far from settled. We hope that our analysis will encourage others to take up the quest, and also to reconsider the related *Chebyshev problem* [3, 7, 17].

## References

- [1] A. Araujo. ‘The nonexistence of smooth demand in general Banach spaces’. *J. Math. Econom.* **17** (4) (1988), 309–319.
- [2] M. Bačák and J. M. Borwein. ‘On difference convexity of locally Lipschitz functions’. *Optimization* **60** (8-9) (2011), 961–978.
- [3] J. M. Borwein. ‘Proximality and Chebyshev sets’. *Optim. Lett.* **1** (1) (2007), 21–32.
- [4] J. M. Borwein and S. Fitzpatrick. ‘Existence of nearest points in Banach spaces’. *Canad. J. Math.* **41** (4) (1989), 702–720.
- [5] J. M. Borwein and A. S. Lewis. *Convex analysis and nonlinear optimization*. CMS Books in Mathematics/Ouvrages de Mathématiques de la SMC, 3 (Springer, New York, 2006), 2nd edn. Theory and examples.
- [6] J. M. Borwein and D. Preiss. ‘A smooth variational principle with applications to subdifferentiability and to differentiability of convex functions’. *Trans. Amer. Math. Soc.* **303** (2) (1987), 517–527.
- [7] J. M. Borwein and J. D. Vanderwerff. *Convex functions: constructions, characterizations and counterexamples*, *Encyclopedia of Mathematics and its Applications*, Volume 109 (Cambridge University Press, Cambridge, 2010).
- [8] J. M. Borwein and Q. J. Zhu. *Techniques of variational analysis*. CMS Books in Mathematics/Ouvrages de Mathématiques de la SMC, 20 (Springer-Verlag, New York, 2005).
- [9] M. Cepedello Boiso. ‘Approximation of Lipschitz functions by  $\Delta$ -convex functions in Banach spaces’. *Israel J. Math.* **106** (1998), 269–284.
- [10] M. Cúth and M. Rmoutil. ‘ $\sigma$ -porosity is separably determined’. *Czechoslovak Math. J.* **63(138)** (1) (2013), 219–234.
- [11] F. S. De Blasi, J. Myjak, and P. L. Papini. ‘Porous sets in best approximation theory’. *J. London Math. Soc.* (2) **44** (1) (1991), 135–142.
- [12] J. Duda. ‘On inverses of  $\delta$ -convex mappings’. *Comment. Math. Univ. Carolin.* **42** (2) (2001), 281–297.
- [13] J. Duda, L. Veselý, and L. Zajíček. ‘On d.c. functions and mappings’. *Atti Sem. Mat. Fis. Univ. Modena* **51** (1) (2003), 111–138.
- [14] Jakub Duda. ‘On the size of the set of points where the metric projection exists’. *Israel J. Math.* **140** (2004), 271–283.
- [15] M. Fabián. ‘Lipschitz smooth points of convex functions and isomorphic characterizations of Hilbert spaces’. *Proc. London Math. Soc.* (3) **51** (1) (1985), 113–126.
- [16] M. Fabián and D. Preiss. ‘On intermediate differentiability of Lipschitz functions on certain Banach spaces’. *Proc. Amer. Math. Soc.* **113** (3) (1991), 733–740.
- [17] J. Fletcher and W. B. Moors. ‘Chebyshev sets’. *J. Aust. Math. Soc.* **98** (2) (2015), 161–231.
- [18] P. Hartman. ‘On functions representable as a difference of convex functions’. *Pacific J. Math.* **9** (1959), 707–713.
- [19] R. Horst, P. M. Pardalos, and N. V. Thoai. *Introduction to global optimization, Nonconvex Optimization and its Applications*, Volume 48 (Kluwer Academic Publishers, Dordrecht, 2000), 2nd edn.
- [20] S. V. Konjagin. ‘Approximation properties of closed sets in Banach spaces and the characterization of strongly convex spaces’. *Dokl. Akad. Nauk SSSR* **251** (2) (1980), 276–280.
- [21] E. Kopecká and J. Malý. ‘Remarks on delta-convex functions’. *Comment. Math. Univ. Carolin.* **31** (3) (1990), 501–510.
- [22] K. S. Lau. ‘Almost Chebyshev subsets in reflexive Banach spaces’. *Indiana Univ. Math. J.* **27** (5) (1978), 791–795.
- [23] J. Lindenstrauss, D. Preiss, and J. Tišer. *Fréchet differentiability of Lipschitz functions and porous sets in Banach spaces*, *Annals of Mathematics Studies*, Volume 179 (Princeton University Press, Princeton, NJ, 2012).
- [24] D. Pavlica. ‘A d.c.  $C^1$  function need not be difference of convex  $C^1$  functions’. *Comment. Math. Univ. Carolin.* **46** (1) (2005), 75–83.

- [25] J.P. Penot. *Calculus without derivatives, Graduate Texts in Mathematics*, Volume 266 (Springer, New York, 2013).
- [26] D. Preiss and L. Zajíček. ‘Fréchet differentiation of convex functions in a Banach space with a separable dual’. *Proc. Amer. Math. Soc.* **91** (2) (1984), 202–204.
- [27] J. P. Revalski and N. V. Zhivkov. ‘Small sets in best approximation theory’. *J. Global Optim.* **50** (1) (2011), 77–91.
- [28] J. P. Revalski and N. V. Zhivkov. ‘Best approximation problems in compactly uniformly rotund spaces’. *J. Convex Anal.* **19** (4) (2012), 1153–1166.
- [29] S. Shalev-Shwartz. ‘Online learning: Theory, algorithms, and applications’, available at <http://www.cs.huji.ac.il/shais/papers/ShalevThesis07.pdf>.
- [30] S. B. Stečkin. ‘Approximation properties of sets in normed linear spaces’. *Rev. Math. Pures Appl.* **8** (1963), 5–18.
- [31] P. T. Thach. ‘D.c. sets, d.c. functions and nonlinear equations’. *Math. Programming* **58** (3, Ser. A) (1993), 415–428.
- [32] P. T. Thach and H. Konno. ‘D.c. representability of closed sets in reflexive Banach spaces and applications to optimization problems’. *J. Optim. Theory Appl.* **91** (1) (1996), 1–22.
- [33] L. Veselý and L. Zajíček. ‘Delta-convex mappings between Banach spaces and applications’. *Dissertationes Math. (Rozprawy Mat.)* **289** (1989), 52.
- [34] L. Veselý and L. Zajíček. ‘On D.C. mappings and differences of convex operators’. *Acta Univ. Carolin. Math. Phys.* **42** (2) (2001), 89–97. 29th Winter School on Abstract Analysis (Lhota nad Rohanovem/Zahrádky u České Lípy, 2001).
- [35] L. Zajíček. ‘Differentiability of the distance function and points of multivaluedness of the metric projection in Banach space’. *Czechoslovak Math. J.* **33(108)** (2) (1983), 292–308.
- [36] L. Zajíček. ‘On  $\sigma$ -porous sets in abstract spaces’. *Abstr. Appl. Anal.* (5) (2005), 509–534.

Jonathan M. Borwein, Centre for Computer-assisted Research Mathematics and its Applications (CARMA), School of Mathematical and Physical Sciences, University of Newcastle, Callaghan, NSW 2308, Australia  
e-mail: [jonathan.borwein@newcastle.edu.au](mailto:jonathan.borwein@newcastle.edu.au)

Ohad Giladi, Centre for Computer-assisted Research Mathematics and its Applications (CARMA), School of Mathematical and Physical Sciences, University of Newcastle, Callaghan, NSW 2308, Australia  
e-mail: [ohad.giladi@newcastle.edu.au](mailto:ohad.giladi@newcastle.edu.au)